Award Number: W81XWH-13-2-0032

TITLE: Increasing Treatment Seeking Among At-Risk Service Members Returning

from Warzones

PRINCIPAL INVESTIGATOR: Tracy Stecker, PhD

CONTRACTING ORGANIZATION: Dartmouth Medical School

Lebanon, NH 03766

REPORT DATE: March 2016

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Affington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE	2. REPORT TYPE	3. DATES COVERED
March 2015	Annual Report	28 Feb 2015 - 27 Feb 2016
4. TITLE AND SUBTITLE		5a. CONTRACT NUMBER
		W81XWH-13-2-0032
Increasing Treatment Seeking Among	5b. GRANT NUMBER	
from Warzones	W81XWH-13-2-0032	
Trom warzones		5c. PROGRAM ELEMENT NUMBER
a AUTHOR(O)		5 L DDO ISOT NUMBER
6. AUTHOR(S)	d Martaglanda	5d. PROJECT NUMBER
Tracy Stecker, PhD Dartmou	th Medical School	F. TACK NUMBER
		5e. TASK NUMBER
		5f. WORK UNIT NUMBER
E-Mail: tracv.stecker@dartmouth.edu		
7. PERFORMING ORGANIZATION NAME Trustees of Dartmouth Col 1 Hinman Dartmouth College	S) AND ADDRESS(ES)	8. PERFORMING ORGANIZATION REPORT NUMBER
Hanover NH 03755-4099		
9. SPONSORING / MONITORING AGENC	Y NAME(S) AND ADDRESS(ES)	10. SPONSOR/MONITOR'S ACRONYM(S)
U.S. Army Medical Research and I	, , , , , , , , , , , , , , , , , , , ,	
Fort Detrick, Maryland 21702-501	2	
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)

12. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for Public Release; Distribution Unlimited

13. SUPPLEMENTARY NOTES

14. ABSTRACT

We propose a randomized controlled clinical trial of 1,200 military service members who are at increased risk for suicide but not currently in behavioral health treatment for the purpose of determining if a brief intervention improves the initiation of treatment. Participants assigned to the treatment condition will be presented an individualized CB intervention. The CB intervention takes 45-60 minutes, is delivered by phone, and has been shown to promote treatment-seeking including in a preliminary study of OEF/OIF Veterans with elevated posttraumatic stress disorder (PTSD) symptoms. Participants will be assessed at baseline and at 1-month, 3-month, and 6-month follow-up. Analyses are based on logistic and mixed effect models.

15. SUBJECT TERMS

Nothing listed

16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U	UU	6	19b. TELEPHONE NUMBER (include area code)

Table of Contents

<u>Page</u>

Introduction	1
Body	2-3
Key Research Accomplishments	n/a
Reportable Outcomes	n/a
Conclusion	n/a
References	n/a
Appendices	n/a

INTRODUCTION TO SPECIFIC AIMS

Reducing suicide is a national priority and an urgent concern within the Department of Defense and the Department of Veterans Affairs. The passage of the Joshua Omvig Veterans Suicide Prevention Act highlights the importance that stakeholders place on developing and implementing a comprehensive program to reduce suicide among U.S. service members and Veterans. Rates of suicide among active duty service members have increased dramatically since 2005, and there is great concern that elevated risk will carry over following discharge from active service. *The goal of this study is to improve initiation of behavioral health (i.e., mental health, substance use) treatment services among untreated, at-risk U.S. military service members.* The goal to facilitate behavioral health treatment is consistent with recommendations provided in reports by the Department of Defense, U.S. Army, U.S. Surgeon General, and the Institute of Medicine.

We are conducting a randomized controlled clinical trial of 1,200 military service members who are at increased risk for suicide but not currently in behavioral health treatment for the purpose of determining if a brief intervention improves attitudes toward behavioral health treatment and initiation of treatment. Advertisements are used to recruit service members who report current suicidal ideation or a history of suicide attempt on standard screening items, suggesting they are at risk for suicide. Participants assigned to the treatment condition are presented an individualized cognitive-behavioral (CB) intervention. The CB intervention takes 45-60 minutes, is delivered by phone, and has been shown to promote treatment-seeking including in a preliminary study of OEF/OIF Veterans with elevated posttraumatic stress disorder (PTSD) symptoms. Participants are assessed at baseline and at 1-month, 3-month, 6-month, and 12-month follow-up. Analyses are based on logistic and mixed effect models. **Specific Aims are as follows**:

- Test the effectiveness of the intervention on attitudes toward behavioral health treatment among at-risk service members.
 <u>Hypothesis 1a</u>: Participants receiving the CB intervention will have significant increases in positive attitudes about treatment at 1-month follow-up compared to controls.
 <u>Hypothesis 1b</u>: Participants receiving the CB intervention will have significant increases in the intention to initiate behavioral health treatment compared to controls.
- 2) Test the effectiveness of the intervention on the initiation of and adherence to behavioral health treatment.
 <u>Hypothesis 2a</u>: Participants receiving the CB intervention will be more likely to initiate behavioral health treatment than participants in control group during 6-month follow-up.
 <u>Hypothesis 2b</u>: Participants receiving the CB intervention will attend more behavioral health treatment sessions than participants in the control group over 6-month follow-up.

The brief (45-60 min), practical (delivered by phone), and promising nature of the CB intervention (favorable preliminary data in OEF/OIF Veterans with elevated PTSD symptoms) indicate its potential for wider implementation to reduce suicide risk among service members.

ACCOMPLISHMENTS

Recruitment for this trial began on March 2, 2013. We aimed to recruit approximately 250 individuals during each year of the trial, and have 748 participants as of March 31, 2016, indicating that we are right on schedule. Of the 748 active participants, 619 of them are separated from the service, 104 are active in the National Guard or Reserves, and 25 are Active Duty.

Recruitment is ongoing and on track.

Baseline characteristics.

Characteristics of the participant sample that have been entered into the data management system are presented in Table 1. The mean age of participants is approximately 31 years old, and 90% of the sample is male. The majority of participants identify their race as white, non-Hispanic. The vast majority reported service in the Army. Baseline symptom severity scores indicate that both the intervention and control groups reported moderately severe symptoms of depression (mean score = 18) as measured by the PHQ9 and PTSD (mean score = 62) as measured by the PCL.

Forty-two percent of the sample reported a previous suicide attempt during the baseline assessment. The majority of the attempts involved the use of medications and or alcohol. Overdosing on medications is also the most frequently reported method when asked if they have a plan for suicide. Other reported methods involve guns, cutting, car accidents, and carbon monoxide poisoning.

We have had four participants die during the trial. One participant died from heart failure related to the use of fentanyl, three died as a result of drug overdose. One of these overdoses was categorized as a suicide. All of these outcomes were reported to all IRB's involved and were judged to be unrelated to study participation.

No SAE's have occurred during this trial period.

REPORTABLE OUTCOMES

No reportable outcomes are available for this trial at this time.

Table 1: Baseline characteristics of the sample

Characteristic	Intervention Participants (n = 362)		Control Participants (n = 369)		
	n	%	n	%	
Male	325	89.8	331	89.7	
Female	37	10.2	38	10.3	
Ethnicity					
White, non-Hispanic	242	66.9	245	66.4	
White, Hispanic	24	6.6	34	9.2	
Black, non-Hispanic	45	12.4	39	10.6	
Black, Hispanic	4	1.2	2	.5	
Native American	4	1.1	8	2.2	
Asian/Pacific Islander	12	3.3	11	3.0	
Other/or mixed race/ethnicity	28	7.7	28	7.6	
Missing	3	.8	2	.5	
Branch of Service					
Army	237	68.4	263	71.3	
Navy	29	8.1	31	8.4	
Airforce	26	7.2	23	6.2	
Marine	54	14.9	51	13.8	
Coast Guard	0	0	1	0.3	
Refused	1	.4	0	0	
Missing	2	.8	0	0	
	Mean	Range	Mean	Range	
Age	31.41	21-58	31.06	21-67	
PHQ-9 score at Baseline	17.74	5-27	18.45	4-27	
PCL score at Baseline	61.96	18-85	62.23	21-85	

 Table 2: Treatment at 1 month follow-up

Characteristic	Intervention	Participants	Control Participants		
	n	%	n	%	
Scheduled appointment	81	33.8	63	25.3	
Attended treatment	53	22.1	24	9.6	
Number of sessions attended	Mean	Range	Mean	Range	
	2.7	0-23	1.97	0-5	